

CLINICAL VIGNETTE

Glenohumeral *Cryptococcus neoformans* septic arthritis: A Case Report

Casimir Dowd, MD¹, Zachary Rubin, MD², Benjamin Bengs, MD¹

¹UCLA, Department of Orthopaedic Surgery, David Geffen School of Medicine
10945 Le Conte Ave, Ste 33-55 PVUB, Los Angeles, CA 90095

²UCLA, Department of Infectious Disease, David Geffen School of Medicine
Box 951720, 12-159 CHS Los Angeles, CA 90095-1720

Correspondence: bbengs@mednet.ucla.edu

Introduction

Cryptococcus neoformans is an encapsulated yeast-like fungus and a human pathogen throughout the world. Ubiquitous in soil and commonly present in bird excreta, *Cryptococcus* primarily enters the body through inhalation¹. Although cryptococcosis can occur in immunocompetent hosts, it is much more commonly found in patients with deficits in immunity such as HIV/AIDS, those with hematologic malignancies or those undergoing treatment with systemic corticosteroids^{2,3}. Infection can result in a primary pulmonary process with a variable presentation from asymptomatic to fulminant pneumonia. More commonly, cryptococcosis results in CNS infection secondary to hematogenous and lymphatic dissemination from the initial pulmonary site of entry and can produce a life-threatening meningoencephalitis^{1,4}. Dissemination can also result in a localized or widespread cutaneous and mucocutaneous infection. Rarely, musculoskeletal infection produces a septic arthritis or osteomyelitis, with bony involvement occurring in about 10% of systemic cryptococcosis^{2,3}. There have been 24 cases of arthritis caused by *Cryptococcus* described in the English language literature⁵⁻⁷. To our knowledge, this is the first reported case of glenohumeral joint septic arthritis due to *Cryptococcus*, and the second reported occurrence of *Cryptococcus* arthritis associated with systemic lupus erythematosus (SLE).

Case Report

The patient is a 44 year-old African American female who presented with one month of right shoulder pain. The patient complained of chills, subjective fevers, and nausea. Medical history

includes idiopathic thrombocytopenic purpura (ITP) for which she underwent splenectomy, and recurrent contralateral *Mycobacterium tuberculosis* axillary lymphadenitis for which she was being treated with rifampin, ethambutol, isoniazid, pyridoxine and pyrazinamide. The patient carried a diagnosis of systemic lupus erythematosus, but was not taking any rheumatologic medications at that time.

Pertinent exam findings were a fever of 101.3F, right shoulder pain with passive and active range of motion, and tenderness and swelling about the shoulder.

Laboratory studies revealed a mildly elevated white blood cell count of 15,900, C-reactive protein of 10.7, and a erythrocyte sedimentation rate of 40.0.

X-ray of the shoulder showed inferior humeral head subluxation and a joint effusion (Figure 1). MRI showed glenohumeral effusion and humeral head changes consistent with septic glenohumeral arthritis and humeral head osteomyelitis (Figure 2). Glenohumeral aspiration revealed gross purulence. Gram's stain revealed yeast forms and aerobic cultures ultimately grew *Cryptococcus neoformans*. Mycobacterial stains and culture, routine cultures and serologic testing for coccidioidomycosis were negative.

In surgery, glenohumeral arthroscopic incision and debridement with open humeral head curettage was performed. Operative findings were consistent with septic arthritis and humeral osteomyelitis.

Infectious disease consultation was obtained and fluconazole 800mg qday was started. HIV testing was negative. Serum cryptococcal antigen was 1 to greater than 2,275. Given the high risk for CNS involvement, a lumbar puncture was performed and was negative for CNS infection. The patient's hospital course was complicated by a right upper extremity deep venous thrombosis and Herpes simplex lesions on her back, both of which resolved with medical treatment. The patient was discharged on fluconazole 800mg daily as well as rifabutin, INH, and pyridoxine for her preexisting tuberculous lymphadenitis.

At 3 months follow-up, she was clinically free from infection and her shoulder tenderness and range of motion had improved significantly.

Discussion

Septic arthritis is a rare complication of cryptococcal infection with only 24 previously reported cases in the literature⁵⁻⁷. The majority of the reported cases involved isolated infections of the knee joint (9), or were polyarticular (7) of which 6 included infections of the knee, making it by far the most commonly afflicted site. There have been several cases involving joints of the upper extremity, including elbow, acromioclavicular, sternoclavicular and wrist joints, however to our knowledge, there have been no previously reported cases involving the glenohumeral joint.

Defects in cell-mediated immunity especially predispose to *Cryptococcus* infection. Of the previous 24 cases, 6 were solid organ transplant recipients, 2 had AIDS, and 3 with sarcoidosis. 9 of 24 patients were on immunosuppressive therapy, most commonly prednisone and azathioprine. Only 4 of 24 patients had no known underlying disease prior to development of cryptococcal arthritis. There has only been one patient previously reported to have SLE as their only predisposing risk factor³.

Although no specific guidelines exist for managing cryptococcal arthritis, adhering to the recommendations for management of cryptococcemia appear to be reasonable in treating joint involvement⁵. Patients with cryptococcal arthritis should be evaluated carefully for disseminated disease and involvement of the central nervous system. Successful management of isolated cryptococcal

arthritis involves initial administration of either amphotericin B or oral fluconazole with or without oral flucytosine for 6-12 months⁸ and often requires surgical intervention (9/24 previous cases refs). Depending on the severity of immunosuppression and persistent or recurrent disease, chronic suppressive azole administration, as is suggested for HIV patients may be considered⁹.

The vast majority of septic arthritis continues to be bacterial. However, atypical pathogens such as *Cryptococcus neoformans* should be considered in the differential, particularly in patients with compromised cellular immunity.

FIGURE LEGENDS

Image 1 – Anteroposterior(AP) Xray of right shoulder demonstrating inferior subluxation of humeral head consistent with joint effusion



Image 2 -- Sagittal T2 Weighted MR demonstrating diffuse soft tissue infiltration and proximal humeral fluid collection consistent with osteomyelitis



REFERENCES:

1. **Murray PR, Rosenthal KS and Pfaller MA.** Cryptococcosis. In: Medical Microbiology, 5th edition. Philadelphia: Elsevier-Mosby; 2005. p.787-89.
2. **Chayakulkeeree M, Perfect JR.** Cryptococcosis. *Infect Dis Clin North Am.* 2006 Sep;20(3):507-44, v-vi. Review. PubMed PMID: 16984867.
3. **Perfect JR, Durack DT, Gallis HA.** Cryptococemia. *Medicine* (Baltimore). 1983 Mar;62(2):98-109. PubMed PMID: 6338344.
4. **Perfect JR.** Cryptococcus neoformans. In: Mandell, Bennet and Dolin: Principles and Practices of Infectious Diseases, 6th edition. Philadelphia: Churchill Livingstone; 2005. p. 2997-3009.
5. **Bruno KM, Farhoomand L, Libman BS, Pappas CN, Landry FJ.** Cryptococcal arthritis, tendinitis, tenosynovitis, and carpal tunnel syndrome: report of a case and review of the literature. *Arthritis Rheum.* 2002 Feb;47(1):104-8. Review. PubMed PMID: 11932886.
6. **McDonough EB Jr, Krishnamurthy AB.** Cryptococcal arthritis of the knee in an immunocompetent host. *Am J Orthop* (Belle Mead NJ). 2005 Mar;34(3):127-8. PubMed PMID: 15828515.
7. **Ortiz M, Gonzalez E, Munoz MA, Andres A.** Cryptococcal monoarthritis without systemic involvement in a renal transplant patient. *Transplantation.* 2004 Jul 27;78(2):301-2. PubMed PMID: 15280697.
8. **Saag MS, Graybill RJ, Larsen RA, Pappas PG, Perfect JR, Powderly WG, Sobel JD, Dismukes WE.** Practice guidelines for the management of cryptococcal disease. Infectious Diseases Society of America. *Clin Infect Dis.* 2000 Apr;30(4):710-8. Epub 2000 Apr 20. PubMed PMID: 10770733.
9. **Chen HS, Tsai WP, Leu HS, Ho HH, Liou LB.** Invasive fungal infection in systemic lupus erythematosus: an analysis of 15 cases and a literature review. *Rheumatology* (Oxford). 2007 Mar;46(3):539-44. Epub 2006 Oct 13. Review. PubMed PMID: 17043051.

Submitted on January 14, 2012