

## CLINICAL VIGNETTE

# Right Ventricular Metastasis of Hepatocellular Carcinoma

Vaughn Eyvazian, M.D.<sup>1</sup> and Kamran Shamsa, M.D.<sup>2</sup>

<sup>1</sup>Department of Medicine, Ronald Reagan UCLA Medical Center, Los Angeles, CA

<sup>2</sup>Department of Cardiology, Ronald Reagan UCLA Medical Center, Los Angeles, CA

A 71-year-old female presents for elective cryoablation of a left adrenal metastasis complicated by hypertensive emergency and NSTEMI with incidental discovery of a right ventricular (RV) pedunculated mass. The patient has a past medical history of hepatitis C virus (HCV) cirrhosis complicated by metastatic hepatocellular carcinoma (HCC) requiring multiple RFA, non-obstructive CAD, and hyperlipidemia. Her oncologic history is significant for metastasis to brain, lung, and left adrenal requiring Y-90 radioembolization, cryoablation, and systemic sorafenib therapy. Upon cardiology follow-up, a new RV pedunculated mass was discovered on TTE (Figure 1). Initial differential diagnosis included HCC metastasis, infective endocarditis, and benign tumor. At this time, she reported mild progressive dyspnea but denied any other new heart failure symptoms. Pertinent physical exam findings include worsening 2/6 LLSB systolic ejection murmur, lack of RV heave or PMI displacement, JVP of 6 cm, absence of crackles and lower extremity edema, Temperature 35.8°C (96.4°F), and BP 120/78. Labs were notable for ESR 43 mm/hr, CRP 0.4 mg/dL with negative bacterial and fungal blood cultures.

Advanced imaging with cardiac MRI showed a 5.8 x 3.5 cm lobular mass along the RV free wall with a large intraluminal component and small polypoid component prolapsing through the tricuspid valve (Figure 2). There was interval increase from previous chest CT imaging. The pedunculated mass showed evidence of narrowing of the RV outflow tract with flow exenteration in the infundibulum. Diagnosis of cardiac metastasis was confirmed with high suspicion. In addition, a right upper lobe pulmonary artery embolus secondary to right ventricular mass was identified. Eventually the patient succumbed to metastatic disease.

### Discussion

Intracardiac metastases are infrequent, and discontinuous ventricular extension is exceedingly rare.<sup>1</sup> In metastatic HCC, involvement is usually isolated to the lung, peritoneum, bone, or adrenals.<sup>1</sup> Most cardiac metastases affect the right atrium and originate with renal cell carcinoma, melanoma, thyroid carcinoma, or testicular carcinoma.<sup>1,2</sup> HCC carries particular affinity for hematogenous spread and vascular extension.<sup>1</sup> Review of 81 Nigerian patients with HCC showed that 44% of patients had post-mortem evidence of tumor thrombi or metastasis to the heart.<sup>3</sup> Additional post-mortem evaluation suggest 2.7-4.1% of patients with metastatic HCC have atrial involvement.<sup>4,5</sup> Overall cardiovascular incidence may be


underestimated; however, development of new cardiovascular symptoms in advanced HCC may necessitate further imaging with echocardiography or cardiac MRI.<sup>5</sup> Use of a multidisciplinary treatment approach including radiation, systemic chemotherapy, and surgical removal of cardiac metastasis may provide clinical benefit but must be used on a case by case basis.

### Figures

**Figure 1.** Transthoracic echocardiogram with a pedunculated mass traversing the tricuspid valve.



**Figure 2.** Cardiac MRI with lobular mass along the RV free wall.



## REFERENCES

1. **Masci G, Magagnoli M, Grimaldi A, Covini G, Carnaghi C, Rimassa L, Santoro A.** Metastasis of hepatocellular carcinoma to the heart: a case report and review of the literature. *Tumori.* 2004 May-Jun;90(3):345-7. Review. PubMed PMID: 15315319.
2. **Masaki N, Hayashi S, Maruyama T, Okabe H, Matsukawa M, Unno J, Maekawa S, Oka T, Tani M, Matsueda K, et al.** Marked clinical improvement in patients with hepatocellular carcinoma by surgical removal of extended tumor mass in right atrium and pulmonary arteries. *Cancer Chemother Pharmacol.* 1994;33 Suppl:S7-11. PubMed PMID: 8137487.
3. **Ayoola EA, Olubuyide IO, Thomas J.** Cardiovascular systemic invasion by hepatocellular carcinoma: incidence and pattern in a west African population. *Afr J Med Med Sci.* 1994 Mar;23(1):61-6. PubMed PMID: 7839948.
4. **Kojiro M, Nakahara H, Sugihara S, Murakami T, Nakashima T, Kawasaki H.** Hepatocellular carcinoma with intra-atrial tumor growth. A clinicopathologic study of 18 autopsy cases. *Arch Pathol Lab Med.* 1984 Dec;108(12):989-92. PubMed PMID: 6095786.
5. **Kawakami M, Koda M, Mandai M, Hosho K, Murawaki Y, Oda W, Hayashi K.** Isolated metastases of hepatocellular carcinoma in the right atrium: Case report and review of the literature. *Oncol Lett.* 2013 May;5(5):1505-1508. PubMed PMID:23760591; PubMed Central PMCID: PMC3678705.

Submitted February 27, 2017