

CLINICAL VIGNETTE

Anesthesia for a Patient With Cutaneous Endometriosis

Fei Zheng-Ward, MD and Elizabeth Tsai, MD

Introduction

Endometriosis is endometrial tissue found outside of the uterine cavity and it often leads to infertility and chronic pain. Depending on the location of the endometriosis, the patient can have various presentations that could muddle the diagnosis and delay treatment as there are reports of endometriosis being found in places such as the gastrointestinal track¹ and cutaneous surgical scar sites.² Cutaneous endometriosis is rare and represents less than 1% of all ectopic endometrium³ and it is classified as primary and secondary with primary being without any prior surgical manipulation and secondary is as a result of a surgical history.

Secondary endometriosis is usually a result of iatrogenic implantation of endometrial cells during surgery and it can be found after open or laparoscopic cases, while the pathogenesis of primary endometriosis is still uncertain.³ Cutaneous endometriosis can very rarely be complicated with a fistula formation, however, these conditions have been treated successfully via surgical resection.⁴

Case Report

A 33-year-old female presented to our preoperative anesthesia clinic for excision of a 1x2 cm cutaneous endometriosis at her cesarean section scar site after her surgery 2 years prior in Venezuela. Her other medical conditions included dysmenorrhea and obesity. The patient stated that the lesion had been there since her cesarean section and that the pain is worse with her menses and it sometimes bleeds with menses as well. She had a CT-guided core biopsy of the lesion which proved to be endometriosis. She had tried oral contraceptive pills and non-steroidal analgesics without success and wanted to proceed to remove the lesion surgically.

On the day of the procedure, the patient was brought to the operating room. Monitors were placed and the patient was given intravenous sedation which included versed, fentanyl and propofol. After she was sedated, she was prepped and draped and was given local anesthetic around the lesion and it was easily removed. The sedation was then turned off at the end of the procedure and the patient was taken to the post anesthesia care unit (PACU). She recovered well and was sent home shortly after that.

Discussion

Endometrial tissue in a surgical scar is not common and can often be misdiagnosed.² In darker skin individuals, cutaneous endometriosis can be mistaken for keloids.⁵ There is a prior report of a woman developing cutaneous endometriosis after caesarean section,⁶ similar to our patient described above. There is another report of a tubocutaneous fistula developing in a post cesarean section surgical scar. Surgery excision was the treatment of choice.⁷ As hormonal therapies can improve symptoms but they do not remove these lesions. Complete wide excision of cesarean scar endometriosis is both diagnostic and therapeutic.⁸

REFERENCES

1. **Tomiguchi J, Miyamoto H, Ozono K, Gushima R, Shono T, Naoe H, Tanaka M, Baba H, Katabuchi H, Sasaki Y.** Preoperative Diagnosis of Intestinal Endometriosis by Magnifying Colonoscopy and Target Biopsy. *Case Rep Gastroenterol.* 2017 Aug 23;11(2):494-499. doi: 10.1159/000475751. eCollection 2017 May-Aug. PubMed PMID: 29033768; PubMed Central PMCID: PMC5624264.
2. **Petrosellini C, Abdalla S, Oke T.** The Many Guises of Endometriosis: Giant Abdominal Wall Endometriosis Masquerading as An Incisional Hernia. *Int J Fertil Steril.* 2018 Jan;11(4):321-325. doi: 10.22074/ijfs.2018.5126. Epub 2017 Oct 12. PubMed PMID: 29043711; PubMed Central PMCID: PMC5641467.
3. **Loh SH, Lew BL, Sim WY.** Primary Cutaneous Endometriosis of Umbilicus. *Ann Dermatol.* 2017 Oct;29(5):621-625. doi: 10.5021/ad.2017.29.5.621. Epub 2017 Aug 25. PubMed PMID: 28966521; PubMed Central PMCID: PMC5597658.
4. **Juneja SK, Tandon P, Chopra I.** Successful pregnancy after excision of cesarean scar endometriosis with utero-vesicocutaneous fistula: A rare case report. *Int J Appl Basic Med Res.* 2016 Oct-Dec;6(4):300-302. PubMed PMID: 27857904; PubMed Central PMCID: PMC5108113.
5. **Kourouma HS, Ecra EJ, Allou AS, Kouyaté M, Kouassi YI, Kloga M, Kouassi KA, Kassi K, Kouamé K, Ahogo C, Gbery IP, Sangaré A.** [Umbilical endometriosis mimicking a keloid in a young black woman: A case report]. *Ann Dermatol Venereol.* 2017 Oct;144(10):617-620. doi: 10.1016/j.annder.2017.05.004. Epub 2017 Jun 23. French. PubMed PMID: 28648517.

6. **Russo VA, Alikhan A.** Painful nodule in the caesarean section scar of a young woman. *Dermatol Online J.* 2015 Nov 18;21(11). pii: 13030/qt4rn7q88d. PubMed PMID: 26632937.
7. **Lopes EN, Damásio LC, Passos LS.** Tubocutaneous Fistula due to Endometriosis - A Differential Diagnosis in Cutaneous Fistulas with Cyclic Secretion. *Rev Bras Ginecol Obstet.* 2017 Jan;39(1):31-34. doi: 10.1055/s-0036-1597754. Epub 2017 Feb 28. PubMed PMID: 28245509.
8. **Uçar MG, Şanlıkan F, Göçmen A.** Surgical Treatment of Scar Endometriosis Following Cesarean Section, a Series of 12 Cases. *Indian J Surg.* 2015 Dec;77(Suppl 2):682-6. doi: 10.1007/s12262-013-0978-1. Epub 2013 Sep 26. PubMed PMID: 26730088; PubMed Central PMCID: PMC4692860.

Submitted May 15, 2018