

CLINICAL VIGNETTE

Evaluation of Vitamin D Levels in Patients With Coccidioidomycosis, a Case Control Study

John Dickey, M.D.¹, Arash Heidari, M.D.², Greti Petersen, M.D.¹,
Brian Jean Ph.D.³, Royce Johnson, M.D.²

(1)Internal Medicine, Kern Medical Center/UCLA, Bakersfield, CA, (2)Infectious Diseases, Kern Medical Center/UCLA,
Bakersfield, CA, (3) Professor of Mathematics Taft College

Introduction

Infection by *coccidioides spp.*, a fungus, found only in the western hemisphere has been a growing concern in endemic areas of Southwest America^{1,2}. Ethnicity, pregnancy, and cell-mediated immunodeficiency have shown to be risk factors for this disease^{3,4}. Immune deficiency plays an important role by increasing the severity of this infection³.

Studies have shown vitamin D deficiency is associated with more severe granulomatous diseases such as tuberculosis⁵⁻⁷. In a recent study, Vitamin D deficiency was associated with severity of liver disease in Human Immunodeficiency Virus (HIV) and Hepatitis C co-infected patients⁸. Vitamin D deficiency has also been linked to worsening oral candidiasis in HIV patients attributed to the down-regulation of calprotectin which influences neutrophil function⁹.

Vitamin D is an important factor in the immune response and affects vitamin D receptors (VDRs) on macrophages, dendritic cells, and activated T and B lymphocytes. The VDRs promote the immune reaction as well as up regulation of the innate immune response such as stimulating defensin and cathelicidin-an antimicrobial peptide^{10,11}.

Vitamin D deficiency has been shown to also inhibit the adaptive immune reaction by inhibiting maturation of the dendritic cells antigen presentation and decreasing T-Cell proliferation^{10,12}. When *coccidioides spp.* spores are inhaled, the natural immune response occurs in two phases. The first response involves neutrophils, monocytes, eosinophils, and natural killer cells. The second phase is composed of macrophages and dendritic cells¹³.

Vitamin D binding protein, known as gc-globulin, is a protein related to the albumin family. This protein transports vitamin D to its target tissue and is found

in plasma, ascitic fluid, cerebrospinal fluid, and on the surface of many proteins. Vitamin D binding protein binds 85-90% of the total 25-Hydroxyvitamin D in circulation. The non-vitamin D-binding protein fraction consists of albumin bound vitamin D, which is the bioavailable form for 10-15% of the total. Less than 1% of vitamin D is in the free form. From prior research, vitamin D-binding protein seems to inhibit actions of vitamin D, because the bound fraction may be unavailable to act on target cells. Genetic variation affects the affinity of the binding protein to vitamin D. The clinical assays measure total vitamin D and do not distinguish fractions bound to carrier proteins¹⁴.

Even though vitamin D deficiency has shown to decrease immune response to tuberculosis and HIV, to our knowledge, no data have been published on vitamin D deficiency and its association with *coccidioidomycosis*.

We compare vitamin D levels in patients with and without coccidioidomycosis in a teaching county hospital population in the San Joaquin Valley.

Methods

Study design:

This is a retrospective, matched case-control study. The patients were older than 18 years and were selected from the Medicine Department clinic of Kern Medical Center from 1990 to 2010. A case was defined as positive for *coccidioidomycosis* either by serology (97%) or by pathology or cultures (31%) (N=118). Controls were patients without any clinical or laboratory findings suggestive of *coccidioidomycosis* (N=472). All cases and controls had vitamin D levels measured. Cases and controls were matched (ratio of 1 to 4) by age, race, and sex and evaluated for co-morbidities. All vitamin D

levels were measured using the Abbott AxSYM Immunoassay System.

Both cases and controls were gender matched with 63% male and 37% females. Both had 16.9% African Americans, 14.4% Caucasians, and 68.6% Hispanics. Among cases 35% of patients had pulmonary *coccidioidomycosis* and 65% with disseminated form with mean age of the cases at 42.8 years old (18-88).

Definition:

The definitions of vitamin D deficiency were similar to previous studies^{15,16}. Vitamin D deficiency was vitamin D OH total <20 ng/ml and Vitamin D insufficiency was a value ≥ 20 and <30 ng/ml^{15,16}. Diabetes Mellitus (DM) was defined as hemoglobin A1C greater than 6.5, previous history of diabetes, or two fasting glucose greater than 126 mg/dl¹⁷. Chronic kidney Disease (CKD) was defined as presence of kidney damage or decreased kidney function for three or more months, irrespective of cause according to the Kidney Disease Outcomes Quality Initiative (KDOQI) and Kidney Disease Improving Global Outcomes (KDIGO) guidelines¹⁸.

Data analysis and statistical methods:

JMP (SAS) version 9 was used for data analysis. Two sample independent t-tests were used to test for significant differences. Odds ratios were used to compare prevalence between case and control groups. Descriptive statistics were also generated for groups of interest. All p-values < 0.05 were considered statistically significant.

Results

The average vitamin D level in cases was 25.9 ng/ml and in controls was 22.5 ng/ml, which was significantly lower than cases $p < 0.01$ (Figure 1).

A numerical analysis of the values showed the vitamin D average for controls was statistically lower than the cases $p < 0.002$. Although averages showed a statistical difference, the prevalence of vitamin D insufficiency (20 < vitamin D < 30) was not statistically higher among cases, OR = 1.52 {CI95 (0.92, 2.5)} $p = 0.0504$.

The prevalence of vitamin D deficiency (<20) was statistically higher in controls OR = 1.88 {CI95(1.1054, 3.1922)} $p = 0.0099$ (Figure 2).

When combining vitamin D deficiency and insufficiency (vitamin D < 30) there was no statistical

difference between the cases and the controls with OR=0.89 {CI95 (0.57,1.39)} $p = 0.30$ (Figure 3).

In terms of comorbidities, 46% of the cases had DM and 7% had CKD. In the control patients, 68% had DM, 23% had CKD, 30% had no co-morbidities, and 6% had other disease (Table 1).

Further analysis found a high prevalence of vitamin D insufficiency and deficiency combined (vitamin D <30) in subjects with DM with an OR=1.766 {CI95 (1.203, 2.593)} $p = 0.0037$, as well as those subjects with CKD with an OR = 1.83 {CI95 (1.106, 3.027)} $p = 0.0187$ (Figures 4 and 5).

We also found the prevalence of DM and CKD were statistically higher in controls compared to cases respectively with an OR = 1.799 {CI95 (1.171, 2.763)} $p = 0.004$ (Figure 6), and an OR=3.789{CI95(1.895, 7.723)} $p = 0.0001$ (Figure 7).

After exclusion of DM and CKD from cases and controls, we found no statistical difference between cases and controls for prevalence of vitamin D deficiency and insufficiency combined, OR = 0.95 {CI95(0.60,1.51)} $p = 0.4241$ (Figure 8,9).

For insufficiency (20 < vitamin D < 30) OR 1.54 {CI95(0.9238, 2.5770)} $p = 0.0487$ the study found a higher prevalence among the cases. For deficiency (vitamin D < 20) the study showed OR = 0.5771 {CI95 (0.3319,1.0038)} $p = 0.0258$ indicating a higher prevalence among the controls.

Discussion

This study showed no correlation between vitamin D levels in patients with or without *coccidioidomycosis*. A recent abstract by University of California Davis investigators, presented at the Annual Infectious Diseases Society of America in 2011 showed similar results¹⁹. These findings are in contrast to the role of vitamin D in tuberculosis, severe liver disease, in coinfecting HIV/HCV, and oral candidiasis in HIV patients.

Our patient population was mostly Hispanic which may not represent the general population.

Furthermore, this study does not explore the effects of free vitamin D versus the bound form. Glucocorticoid or stress hormones are known to decrease vitamin D receptors gene expression in most tissues, which in turn can down regulate metabolic pathways as well as the immune response. Genetic variation of affinity among races for binding protein

to vitamin D has not been studied and could affect result of this study with mostly Hispanic population.

Table 1: Demographics

	Cases (118)	Controls (472)
Age, Average years	42.6	44.5
18 to 39, % (n)	35.6% (42)	35.6% (168)
40 to 59, % (n)	65% (65)	59% (280)
≥60, % (n)	9.3% (11)	9.3% (44)
Gender, % male (n)	63% (75)	63% (300)
Race, % (n)		
Caucasian	14.4% (17)	14.4% (68)
Hispanic	68.6% (81)	68.6% (324)
African American	16.9% (20)	16.9% (80)
Comorbidities, % (n)		
Chronic Kidney Diseases	7%(9)	23%(111)
Diabetes	46%(37)	68%(322)
HIV+	0%(0)	<0.1%(2)
Gastric Bypass	0%(0)	<0.1%(4)
Malignancy	1%(2)	5%(25)
End Stage Liver Disease	1%(2)	1%(6)
Coccidioidomycosis infection % (n)		
Pulmonary	35.5% (42)	0%
Disseminated	64.5% (76)	0%
Any positive culture for <i>C. immitis</i>	31% (37)	0%
Positive complement fixation titers	97.4% (110)	0%




Figure 1: Case or Control versus Vitamin D levels at Baseline




Figure 2: Case or Controls




Figure 3: Case or Controls

In conclusion, this study does not support routine measurement of vitamin D levels in patients with *coccidioidomycosis*.




Figure 4: Comorbidities: DM (Diabetes Mellitus)




Figure 5: Comorbidities: CKD (Chronic Kidney Disease)




Figure 6: Comorbidities: DM (Diabetes Mellitus) in Case and Controls




Figure 7: Comorbidities: CKD (Chronic Kidney Disease) in Case and Controls




Figure 8: DM (Diabetes Mellitus) and CKD (Chronic Kidney Disease) Excluded




Figure 9

REFERENCES

- Centers for Disease Control and Prevention (CDC). Increase in reported coccidioidomycosis--United States, 1998-2011. *MMWR Morb Mortal Wkly Rep.* 2013 Mar 29;62(12):217-21. PubMed PMID: 23535687.
- Pappagianis D; Coccidioidomycosis Serology Laboratory. Coccidioidomycosis in California state correctional institutions. *Ann N Y Acad Sci.* 2007 Sep;1111:103-11. Epub 2007 Mar 1. Review. PubMed PMID: 17332089.
- Woods CW, McRill C, Plikaytis BD, Rosenstein NE, Mosley D, Boyd D, England B, Perkins BA, Ampel NM, Hajjeh RA. Coccidioidomycosis in human immunodeficiency virus-infected persons in Arizona, 1994-1997: incidence, risk factors, and prevention. *J Infect Dis.* 2000 Apr;181(4):1428-34. Epub 2000 Apr 7. PubMed PMID: 10753734.
- Galgiani JN, Ampel NM, Blair JE, Catanzaro A, Johnson RH, Stevens DA, Williams PL; Infectious Diseases Society of America. Coccidioidomycosis. *Clin Infect Dis.* 2005 Nov 1;41(9):1217-23. Epub 2005 Sep 20. Review. PubMed PMID: 16206093.
- Rook GA. The role of vitamin D in tuberculosis. *Am Rev Respir Dis.* 1988 Oct;138(4):768-70. PubMed PMID: 2849343.
- Nnoaham KE, Clarke A. Low serum vitamin D levels and tuberculosis: a systematic review and meta-analysis. *Int J Epidemiol.* 2008 Feb;37(1):113-9. doi: 10.1093/ije/dym247. Review. PubMed PMID: 18245055.
- Talat N, Perry S, Parsonnet J, Dawood G, Hussai R. Vitamin D deficiency and tuberculosis progression. *Emerg Infect Dis.* 2010 May. <http://wwwnc.cdc.gov/eid/article/16/5/09-1693>

- Guzmán-Fulgencio M, García-Álvarez M, Berenguer J, Jiménez-Sousa MÁ, Cosín J, Pineda-Tenor D, Carrero A, Aldámiz T, Alvarez E, López JC, Resino S. Vitamin D deficiency is associated with severity of liver disease in HIV/HCV coinfecting patients. *J Infect.* 2014 Feb;68(2):176-84. doi: 10.1016/j.jinf.2013.10.011. Epub 2013 Nov 1. PubMed PMID: 24184809.
- Sroussi HY, Burke-Miller J, French AL, Adeyemi OM, Weber KM, Lu Y, Cohen M. Association among vitamin D, oral candidiasis, and calprotectinemia in HIV. *J Dent Res.* 2012 Jul;91(7):666-70. doi: 10.1177/0022034512446342. Epub 2012 Apr 25. PubMed PMID: 22538413; PubMed Central PMCID: PMC3383847.
- Bikle D. Vitamin D and immune function. *US Nephrology.* 2009;4(2):27-32.
- Chen S, Sims GP, Chen XX, Gu YY, Chen S, Lipsky PE. Modulatory effects of 1,25-dihydroxyvitamin D3 on human B cell differentiation. *J Immunol.* 2007 Aug 1;179(3):1634-47. PubMed PMID: 17641030.
- Sigmundsdottir H, Pan J, Debes GF, Alt C, Habtezion A, Soler D, Butcher EC. DCs metabolize sunlight-induced vitamin D3 to 'program' T cell attraction to the epidermal chemokine CCL27. *Nat Immunol.* 2007 Mar;8(3):285-93. Epub 2007 Jan 28. PubMed PMID: 17259988.
- DiCaudo DJ. Coccidioidomycosis: a review and update. *J Am Acad Dermatol.* 2006 Dec;55(6):929-42; quiz 943-5. Review. PubMed PMID: 17110216.
- Powe CE, Evans MK, Wenger J, Zonderman AB, Berg AH, Nalls M, Tamez H, Zhang D, Bhan I, Karumanchi SA, Powe NR, Thadhani R. Vitamin D-binding protein and vitamin D status of black Americans and white Americans. *N Engl J Med.* 2013 Nov 21;369(21):1991-2000. doi: 10.1056/NEJMoa1306357. PubMed PMID: 24256378; PubMed Central PMCID: PMC4030388.
- Dawson-Hughes B, Harris SS, Krall EA, Dallal GE. Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N Engl J Med.* 1997 Sep 4;337(10):670-6. PubMed PMID: 9278463.
- Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, Nicholson GC. Annual high-dose oral vitamin D and falls and fractures in older women: a randomized controlled trial. *JAMA.* 2010 May 12;303(18):1815-22. doi: 10.1001/jama.2010.594. Erratum in: *JAMA.* 2010 Jun 16;303(23):2357. PubMed PMID: 20460620.
- American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care.* 2010 Jan;33 Suppl 1:S62-9. doi: 10.2337/dc10-S062. Erratum in: *Diabetes Care.* 2010 Apr;33(4):e57. PubMed PMID: 20042775; PubMed Central PMCID: PMC2797383.

Submitted on May 12, 2014